

MUSCULOSKELETAL SYSTEM:

(THE SKULL):



Ctanial Romes: 8


Fascial Bones: 14

Bones of the Skull

The cranium consists of the following bones, two

of which are paired

- ✓ Frontal bone: 1
- ✓ Parietal bones: 2 ✓
- ✓ Occipital bone: 1
- ✓ Temporal bones: 2
- ✓ Sphenoid bone: 1
- ✓ Ethmoid bone: 1





Bones of the Skull


The facial bones consist of the following, two of which are single:

- ✓ Zygomatic bones: 2 ✓
- ✓ Maxillae: 2 ✓
- ✓ Nasal bones: 2 ✓
- ✓ Lacrimal bones: 2 ✓
- ✓ Vomer: 1 ✓
- ✓ Palatine bones: 2 🔮
- ✓ Inferior conchae: 2 ✓
- ✓ Mandible: 1 ✓

Dr. Aiman Al -Maathidy Sunday 23 February 2025







* Anterior View of the skull:

1 Frontal Lobe:


* orbital cavity: ?

Superiorly: Prontal Bone


Inferiorly: Marillary Bong

Laterally: Zygomatic

Melinity: Process of Frontal + maxillary




2 Nasal Bore:



External Views of the Skull

Anterior View of the Skull


- **❖ The frontal bone**, or forehead bone, curves downward to make the upper margins of the orbits
- *The superciliary arches and the supraorbital notch, or foramen, can be recognized.
- **The orbital margins are bounded by:**
 - √ The frontal bone superiorly,
- √ The zygomatic bone laterally,
- √ The maxilla inferiorly,
- √ The processes of the maxilla and frontal bone medially.



Anterior View of the Skull

❖Within the frontal bone, just above the orbital margins, are two hollow spaces lined with mucous membrane called the frontal air sinuses.


These communicate with the nose and serve as voice resonators.



Anterior View of the Skull


❖ The two nasal bones form the bridge of the nose. Their lower borders, with the maxillae, make the anterior nasal aperture.


The nasal cavity is divided into two by the bony nasal septum, which is largely formed by The Vomer.



13 Maxillary Rones:

Forms:






Anterior View of the Skull

*The two maxillae form the upper jaw, the anterior part of the hard palate, part of the lateral walls of the nasal cavities, and part of the floors of the

orbital cavities.


❖The two bones meet in the midline at the intermaxillary suture and form the lower margin of the nasal aperture.




Dr. Aiman Al -Maathidy
Sunday 23 February 2025


Anterior View of the Skull


- *Below the orbit, the maxilla is perforated by the infraorbital foramen.
- Within each maxilla is a large, pyramid-shaped cavity lined with mucous membrane called the maxillary sinus





Y Zygomalic Bone:






Anterior View of the Skull


❖ The Zygomatic bone forms the prominence of the cheek and part of the lateral wall and floor of the orbital cavity.


❖ The Zygomatic bone is perforated by two foramina for the zygomaticofacial and zygomaticotemporal nerves

*The mandible, or lower jaw, consists of a horizontal body and two vertical rami











- ✓ The frontal bone forms the anterior part of the side of the skull and articulates with the parietal bone at the coronal suture
- ✓ The parietal bones form the sides and roof of the cranium and articulate
 with each other in the midline at the sagittal suture.

√ The parietal bones articulate
with the occipital bone behind,
at the lambdoid suture.





* pail of Temporal Bone:




*The skull is completed at the side by the squamous part of the occipital bone; parts of the temporal bone, namely,


The squamous, tympanic, mastoid process, styloid process, and zygomatic process; and the greater wing of the sphenoid.





The ramus and body of the mandible lie inferiorly









□Note that the thinnest part of the lateral wall of the skull is where the anteroinferior corner of the parietal bone articulates with the greater wing of the sphenoid; this point is referred to as the pterion.

□Clinically, the pterion is an important area because it overlies the anterior division of the middle meningeal artery and vein.





*The infratemporal fossa lies below the infratemporal crest on the greater wing of the sphenoid.

*The pterygomaxillary fissure is a (vertical fissure) that lies within the fossa between the pterygoid process of the sphenoid bone and back of the maxilla. It leads medially into the pterygopalatine fossa

Identify the boundaries of the infratemporal fossa. Temporal Fossa Parietal Frontal Temporal Maxilla Boundaries: Lateral=Ramus of Mandible Lat. Pterygoid Plate Anterior=Maxilla Medial=Lat. Pterygoid Plate Pterygomaxillary Fissure Roof=Sphenoid Infratemporal Fossa

Posterior View of the Skull


✓ The posterior parts of the two parietal bones with the intervening sagittal suture are seen above. Below, the parietal bones articulate with the squamous part of the occipital

bone at the lambdoid suture.

✓On each side the occipital bone articulates with the temporal bone.

✓ In the midline of the occipital bone is a roughened elevation called the external occipital protuberance, which gives attachment to muscles and the ligamentum nuchae.

✓ On either side of the protuberance the superior nuchal lines extend laterally toward the temporal bone





Superior View of the Skull


- >Anteriorly, the frontal bone articulates with the two parietal bones at the coronal suture.
- >Occasionally, the two halves of the frontal bone fail to fuse, leaving a midline

metopic suture. V: 2 Haives of Irontal Bone due to Palive of Pusion

> Behind, the two parietal bones articulate in the midline at the sagittal suture









- ✓ Is the largest and strongest bone of the face
- ✓ It articulates with the skull at the temporomandibular joint. (Mobile)
- ✓ Consists of a horseshoe-shaped body and a pair of rami.
- ✓ The body of the mandible meets the ramus on each side at the angle of


the mandible





- ✓ On the external surface in the midline of the body of the mandible, has a faint ridge indicating the line of fusion of the two halves during development at the symphysis menti.
- ✓ The mental foramen can be seen below the second premolar tooth; it transmits the terminal branches of the inferior alveolar nerve and

vessels.






✓ On the medial surface of the body of the mandible in the median plane are seen the mental spines.

✓ The mylohyoid line can be seen as an oblique ridge that runs backward and laterally from the area of the mental spines to an area below and behind the third molar tooth.

Dr. Aiman Al -Maathidy Sunday 23 February 2025




✓ The submandibular fossa, for the superficial part of the submandibular salivary gland, lies below the posterior part of the mylohyoid line.

✓ The sublingual fossa, for the sublingual gland, lies above the anterior

part of the mylohyoid line.


✓ The upper border of the body
of the mandible is called the
alveolar part; in the adult, it
contains 16 sockets for the
roots of the teeth



- ✓ The lower border of the body of the mandible is called the base.
- ✓ The digastric fossa is a small, roughened depression on the base, on
 either side of the symphysis menti for the anterior bellies of the digastric


muscles attachment.






✓ The ramus of the mandible is vertically placed and has an anterior coronoid process and a posterior condylar process, or head; the two processes are separated by the mandibular notch







- ✓ On the lateral surface of the ramus are markings for the attachment of the masseter muscle.
- ✓ On the medial surface is the mandibular foramen for the inferior alveolar nerve and vessels







- ✓ In front of the foramen is a projection of bone, called the lingula, for the
- ✓ The foramen leads into the mandibular canal, which opens on the lateral surface of the body of the mandible at the mental foramen





✓ The incisive canal is a continuation forward of the mandibular canal beyond the mental foramen and below the incisor teeth.





Sunday 23 February 2025

Dr. Aiman Al -Maathidy

